

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE March 21, 1997		3. REPORT TYPE AND DATES COVERED	
4. TITLE AND SUBTITLE Geostatistical and Neural-Net Seafloor Classification at High Resolution and Related Scaling Properties (part of OPR ARSRP data analysis)				5. FUNDING NUMBERS N00014-95-1-0447	
6. AUTHOR(S) Ute C. Herzfeld					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Institute of Arctic and Alpine Research University of Colorado Boulder, CO 80309-0450				8. PERFORMING ORGANIZATION REPORT NUMBER 1530702	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research, Code 3210A Dr. H. Badley, E. Estalote, T. Travis, B. Kuhn 800 N. Quincy St. ARLINGTON, VA 22217-5660				10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES					
12a. DISTRIBUTION / AVAILABILITY STATEMENT <div style="border: 1px solid black; padding: 5px; text-align: center;"> DISTRIBUTION STATEMENT A Approved for public release; Distribution Unlimited </div>				12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) <div style="font-size: 2em; text-align: center; margin-top: 20px;">19970415 086</div> <div style="text-align: right; margin-top: 20px;">DTIC QUALITY INSPECTED 2</div>					
14. SUBJECT TERMS Seafloor Classification, Geostatistics, Scaling Problems, High-resolution bathymetric data, Western Flank of Mid-Atlantic Ridge				15. NUMBER OF PAGES	
				16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT		

(A) Header Information

Principal Investigator: Dr. Ute C. Herzfeld

Title of Grant: Geostatistical and Neural-Net Seafloor Classification at High Resolution and Related Scaling Properties

Final Report March 21, 1997

Category of Research: ARSRP Surface (Seafloor)

ONR N00014-95-1-0447

Address: Institute of Arctic and Alpine Research, University of Colorado at Boulder, Boulder, CO 80309-0450

e-mail: herzfeld@tryfan.colorado.edu

Phone: (303) 492-6198

Fax: (303) 492-6388

(B) Narrative Documentation

Long-term Research Objectives: My general long-term research objective is the development and application of adequate high-level mathematical techniques for the analysis of spatial data, in particular for remote-sensing data such as acoustic, seismic, bathymetric, and other geophysical data. My interest is in interdisciplinary work in mathematics, measuring techniques, acoustics, environmental information, geology and geophysics.

Scientific Objectives and Report Summary: The objective of my research under this grant is the development of an intelligent system for the automated classification of the seafloor using acoustic data. The work is a contribution to the Acoustical Reverberation Special Research Program (ARSRP). We have developed a method for surface classification, incorporating ideas from the theory of geostatistics (in short, the method has been called "Geostatistical classification method"). Under this project, we have finalized parameter selection and software development for the automated geostatistical seafloor classification method. Thereafter, we applied this method in a geomorphologic segmentation of the Western Flank of the Mid-Atlantic Ridge at 26°North, the area of geophysical survey under the ARSRP in 1992. High-resolution bathymetric data from the 1993 geophysical experiment were also analyzed and compared to the (low-resolution) HYDROSWEET bathymetric data. Surface structures were found to be scale-dependent.

Background and Relationship to Other Research Projects: The problems investigated in this project have arisen in connection with the ARSRP data analysis, but are equally important for the analysis of acoustic and seismic data from deep and shallow ocean environments. Specific questions concern the statistical description of seafloor roughness and the relationship between surface (seafloor) parameters and backscattering properties on the one side and geologic properties on the other side. The results are a contribution

to ocean acoustics, geophysical data analysis, quantitative morphology, and geologic segmentation of the seafloor.

Approach, Accomplishments, and Results:

Geostatistical Seafloor Classification

The objective of automated seafloor classification is to utilize statistical properties in quantitatively characterizing seafloor properties such as roughness and anisotropy. Such spatial characteristics are used to automatically distinguish geological provinces which facilitates a segmentation of the seafloor.

The geostatistical seafloor classification method has been developed in relationship to the analysis of data from the western flank of the Mid-Atlantic Ridge at 26°N. We have analyzed (low-resolution) HYDROSWEEP bathymetric data from the 1992 geophysical experiment in which I participated (cf. Herzfeld 1993) and high-resolution (5 m) data from the 1993 geophysical experiment collected by K. Stewart with the DSL instrument developed at Woods Hole Oceanographic institution (cf. paragraph on "Relationship to other projects").

The geostatistical seafloor classification method is based on calculation of directional variograms as spatial structure functions. Parameters determined from filtered variogram functions are used to compose feature vectors, which have been shown to characterize morphologic prototypes and surface roughness types (Herzfeld 1993a) and therefore facilitate a classification. The variogram is the lag-dependent spatial structure function of geostatistics, the theory of regionalized variables (Journel and Huijbregts 1989). The variograms are filtered, parameters extracted, and a feature vector is composed of the parameters (Herzfeld and Higginson 1996). Discriminants include spacing and strike of abyssal hill terrain, smoothness due to sediment cover, and parameters related to complexity and morphological significance of abyssal hills and their slopes (cf. Figures 1 and 2). Figure 1 shows the MINDIST parameter which characterizes the typical length of features in east-west direction (that is, roughly in the across-strike direction in the survey area on the Western Flank of the Mid-Atlantic Ridge). Figure 2 depicts the parameter P1 indicative of significance of abyssal hill terrain (slope parameter) for the 150km by 100km area shown in Figure 1. More parameters and combinations of parameters are described and applied in Herzfeld and Higginson (1996).

Complications of the automatization concern robustness of parameter estimation, optimal window size, and subselection of data. By moving a classification operation through the study area and color-coding property classes, seafloor classification maps are obtained. The concepts of characteristic parameters, feature vectors, and discrimination criteria are introduced in Herzfeld and Higginson (1996). The updated version of the program contains an option for automatic selection of the locally dominant direction in areas with anisotropies (typical for the ridge flank). This search succeeds in identifying structures inobvious to the eye.

Seafloor Segmentation

The geostatistical classification method has been applied in segmentation of the entire 600 km by 250 km area of the 1992 geophysical survey. The segmentation is based on HYDROSWEEP data. It should be mentioned that the method is robust enough to evaluate data with relatively high error levels and with gaps in the data coverage (both of which occurred in the 1992 HYDROSWEEP data).

Maps of parameters MINDIST, POND, and significance of abyssal hill terrain (slope parameter P1 and relative-size parameter P2) and of parameter combinations have been produced. The map of parameter combination MINDIST and POND is given in Figure 3. The POND parameter shows flat areas, typically




Figure 1.
Parameter HINDIST



Figure 2.
Parameter P1




Figure 3.
Combination of POB and HINDIST parameters.
Basis for segmentation of 600 x 250 km off-axis area.

sediment ponds (notice it depicts the interior of larger ponds due to the moving window operation). Areas without significant features are marked in black (no min-max sequence found). Connected black areas trending from lower right to upper left in the map in Figure 3 delineate the location of discontinuities, which separate three ridge flank segments of different characteristics. The central segment has with distance from the ridge crest (spreading center) gradually increasing sizes of characteristic spacing of abyssal hill terrain, this is indicative of relatively undisturbed Atlantic crust. The northern segment appears to be overprinted by a process other than seafloor spreading, which is obvious from the less parallel and less connected smaller color patches. A third segment can be distinguished in the lower left corner of the map. Some of this interpretation is inferred from maps of other parameters and multidirectional analysis maps not shown in this brief report. The results will appear in Herzfeld (1997).

Results of our work have been presented at the 1994 AGU Fall meeting, at the 1995 meeting of the International Union for Geodesy and Geophysics in Boulder, and at the 1996 meeting of the European Geophysical Society (cf. Abstracts).

Scale-dependent Properties ("Fractal analysis")

Theoretical work on fractals and scaling properties and analysis of seafloor profiles from various parts of the seafloor indicated that the ocean floor does not follow a self-similar or self-affine model (Herzfeld 1993b, Herzfeld et al. 1993a, Herzfeld et al. 1995). Analysis of DSL120 high-resolution data with the classification method revealed once more that the characteristics of the seafloor at 5 m resolution are different from those at about 125 to 200 m resolution. If the seafloor were self-similar, maps of parameter P2 would simply show the same pattern and colors at both resolutions, which is not the case. Rather than just show that features are different, the classification maps yield information on the scale-dependent morphologic characteristics.

Relationship to Other Projects:

The project is part of the ARSRP data analysis. Data have been collected during the cruises EW9208 in 1992 (Chief Scientists B. Tucholke and M. Kleinrock) and during the geophysical cruise in 1993 (Chief Scientists K. Stewart and B. Tucholke). I participated in the 1992 cruise with the Maurice Ewing. Development of our geostatistical classification system benefited from the development of graphics programs by M. Marra and K. Stewart (WHOI). Ken Stewart's group has been working on classification, but on a different scale and using a different approach. We have given a few joint presentations at ARSRP meetings and at the IUGG meeting (cf. Abstract). Results of the geostatistical surface classification can be made available to the research community for testing of acoustic models and to aid in geologic investigations.

References:

- Davis, J.C. and U.C. Herzfeld (eds.), *Computers in geology: 25 years of progress*, Internat. Assoc. Math. Geol. Studies in Mathematical Geology, no. 5, Oxford University Press, New York (1993), 298 pp.
- Herzfeld, U.C., 1993a, A method for seafloor classification using directional variograms, demonstrated for data from the western flank of the Mid-Atlantic Ridge, *Math. Geology*, vol. 25, no. 7, p. 901-924.
- Herzfeld, U.C., 1993b, Fractals in geosciences - challenges and concerns, in: Davis, J.C. and U.C. Herzfeld (eds.), *Computers in geology: 25 years of progress*, Internat. Assoc. Math. Geol. Studies in Mathematical Geology, no. 5, Oxford University Press, New York, 217-230.
- Herzfeld, U.C., 1993c, Application of geostatistical classification, Autumn ARSRP Research Symposium, 1-3 December 1993, La Jolla, CA, Report.
- Herzfeld, U.C., 1997, Geological and morphological provinces of the western Mid-Atlantic Ridge flank at 26°North, derived by automated geostatistical seafloor classification (in prep.).
- Herzfeld, U.C., and C.A. Higginson, 1996, Automated geostatistical seafloor classification - principles, parameters, feature vectors, and discrimination criteria, *Computers & Geosciences*, vol. 22, no. 1, p. 35-52.
- Herzfeld, U.C., I.I. Kim, J.A. Orcutt, and C.G. Fox, 1993a, Fractal geometry and seafloor topography: Theoretical concepts versus data analysis for the Juan de Fuca Ridge and the East Pacific Rise, *Annales Geophysicae*, vol. 11, no. 6, p. 532-541.
- Herzfeld, U.C., M. Marra, and W.K. Stewart, 1993b, A variogram method for seafloor classification, Spring ARSRP Research Symposium, La Jolla, CA, Report.
- Herzfeld, U.C., I.I. Kim, and J.A. Orcutt, 1995, Is the ocean floor a fractal?, *Math. Geology*, vol. 27, no. 3, p. 421-462.
- Journel, A.G., and C. Huijbregts, 1989, *Mining geostatistics*: London, Academic Press, 600 pp.
- Tucholke, B.E., and J. Lin, 1994, A geological model for the structure of ridge segments in slow-spreading ocean crust, *J. Geophys. Res.*, vol. 99, no. B6, p. 11937-11958.

Publications under this project

Reviewed Publications

- (1) HERZFELD, U.C., I.I. KIM, and J.A. ORCUTT, Is the ocean floor a fractal? *Math. Geol.*, vol. 27, no. 3 (1995), p. 421-462
- (2) HERZFELD, U.C. and C.A. HIGGINSON, Automated geostatistical seafloor classification — principles, parameters, feature vectors, and discrimination criteria, *Computers & Geosciences* vol. 22, no. 1 (1996), p. 35-52
- (3) HERZFELD, U.C., Geological and morphological provinces of the western Mid-Atlantic Ridge flank at 26°North, derived by automated geostatistical seafloor classification, *J. Geodynamics* (in preparation)

Abstracts

- (1) HERZFELD, U.C., Automated classification of geologic and geomorphologic provinces of the Western Flank of the Mid Atlantic Ridge, *Eos Trans. Amer. Geophys. Union*, vol. 75, no. 44 suppl. (1994), p. 579 (Amer. Geophys. Union 1994 Fall Meeting, San Francisco, California, USA, Dec. 1994)
- (2) HERZFELD, U.C., C.A. HIGGINSON, K. STEWART and M. MARRA, Segmentation of the Mid-Atlantic Ridge flank at 26°N, derived by automated geostatistical seafloor classification, XXI General Assembly Int. Union Geodesy Geophys., Boulder, Colorado, 2-14 July 1995, Abstracts, Week A, p. A6
- (3) HERZFELD, U.C., Geostatistical methods for interpolation and classification of remote-sensing data, Tagung "Mathematische Methoden in der Geodäsie", Mathematisches Forschungsinstitut Oberwolfach, 1-7 October 1995, Germany
- (4) Segmentation of the western flank of the Mid-Atlantic Ridge at 26°N, an application of automated geostatistical seafloor classification, *Annales Geophysicae*, vol. 14, suppl. 1, part 1 (1996), p. C196 (21st General Assembly European, Geophys. Soc., 6-10 May 1996, Den Haag, The Netherlands)

Statistical Information:

Publication list: enclosed

Graduate Students: C. A. Higginson

Service on Committees:

Associate Editor: Computers & Geosciences, International Association for Mathematical Geology (since 1995, Assistant Editor 1987–1994)

Assistant Editor: Journal of Geodynamics, European Geophysical Society (since 1994)

Assistant Editor: Mathematical Geology, International Association for Mathematical Geology (since 1994)

Report sent cc: J.A. Orcutt, C.A. Higginson, K. Stewart, M. Marra, J. Syvitsky

Ute Christina Herzfeld

RESUME

Born: November 24, 1960, Lieser, Kreis Bernkastel-Wittlich, Germany

Education

- 1978-1983 Scholarship, German National Scholarship Foundation, Bonn, Germany
- 1980-1981 Scholarship for Studies Abroad, German National Scholarship Foundation, Bonn, at the University of Warwick, Coventry, Great Britain
- Dec. 1983 State Examination (Master's Degree), Mathematics, Evangelical Theology, Johannes Gutenberg-Universität Mainz, Germany (summa cum laude)
- 1982-1986 Study of geology, oceanography, geophysics, physical geography, glaciology
- 1984-1986 Doctoral Fellowship, German National Scholarship Foundation, Bonn
- June 1986 Doctoral Degree (Dr. rer. nat.), Mathematics secondary subjects applied mathematics and geosciences, Johannes Gutenberg-Universität Mainz
- 1986 Diploma of Postgraduate Studies, Mathematical Geology, Free University of Berlin, Germany

Positions and Awards

- 1980-1986 Teaching Assistant in mathematics, Universities of Mainz and Warwick, Coventry, Great Britain
- Oct.-Nov.1984 Assistant Research Scientist during Expedition ANTARKTIS III/1, RV POLARSTERN, Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, Germany
- 1986-1987 Research Geomathematician, Department of Geology, Bayerische Julius-Maximilians-Universität, Würzburg
- July-Aug.1987 Visiting Research Scientist, Centre de Géostatistique, Ecole des Mines de Paris, Fontainebleau, France
- 1987-1988 Postdoctoral Research Fellow (DFG) and Lecturer, Mathematical Geology, Free University of Berlin, Germany
- Dec.1987-Apr.1988 Research Scientist, Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, Expedition ANTARKTIS VI/3
- Oct.1988-Sept.1992 Feodor Lynen Research Fellow, Alexander von Humboldt Foundation, Bonn, Germany, at Scripps Institution of Oceanography, University of California at San Diego, USA
- Jun.1991-Nov.1993 Assistant Research Geomathematician (Project Scientist) at Scripps Institution of Oceanography, University of California San Diego
- 1992 President's Prize of the International Association for Mathematical Geology
- Aug.1993-present Research Associate, Institute of Arctic and Alpine Research, University of Colorado, Boulder

PUBLICATION LIST

Reviewed Publications

- (1) On generalized covering subgroups and a characterization of "pronormal", Arch. Math., vol. 41, no. 5 (1983), p. 404-409
- (2) Frattiniklassen und maximale Teilklassen, insbesondere zu Formationen endlicher Gruppen, Dissertation Fachbereich Mathematik, Johannes Gutenberg-Universität Mainz, West Germany, D 77, (1986), 134 p.
- (3) Frattiniklassen and maximal subclasses, in particular of saturated formations of finite groups, J. Algebra, vol. 117, no. 1 (1988), p. 9-10
- (4) Frattiniklassen of formations of finite groups, Boll. Unione Mat. Ital., vol. 7, no. 2-B (1988), p. 601-611
- (5) HERZFELD, U.C. and M. SONDERGARD, MAPCOMP - A FORTRAN 77-program for weighted thematic map comparison, Computers & Geosciences, vol. 14, no. 5 (1988), p. 699-713
- (6) HERZFELD, U.C. and P. HOLMLUND, Geostatistical analyses of radio-echo data from Scharffenbergbotnen, Dronning Maud Land, East Antarctica, Z. Gletscherkunde Glazialgeol., vol. 24, no. 2 (1988), p. 95-110
- (7) A note on programs performing kriging with nonnegative weights, Math. Geol., vol. 21, no. 3 (1989), p. 391-393
- (8) Geostatistical methods for evaluation of SEABEAM bathymetric surveys: Case studies of Wegener Canyon, Antarctica, Marine Geol., vol. 88 (1989), p. 83-95
- (9) Variography of submarine morphology: Problems of deregularization, and cartographical implications, Math. Geol., vol. 21, no. 7 (1989), p. 693-713
- (10) Geostatistical software for evaluation of line survey data, applied to radio-echo soundings in glaciology, in: Hanley, J.T. and D.F. Merriam (eds.), Microcomputer applications in geology II, Computers & Geology, vol. 6, Pergamon Press, New York (1990), p. 119-136
- (11) HERZFELD, U.C. and P. HOLMLUND, Geostatistics in glaciology: Implications of a study of Scharffenbergbotnen, Dronning Maud Land, East Antarctica, Annals Glaciol., vol. 14 (1990), p. 107-110
- (12) Cova functions for unevenly and noncorrespondingly spaced processes, Computers & Geosciences, vol. 16, no. 5 (1990), p. 733-749
- (13) HERZFELD, U.C. and D.F. MERRIAM, A map comparison technique utilizing weighted input parameters, in: Gaál, G. and D.F. Merriam (eds.), Computer applications in resource estimation and assessment for metals and petroleum, Computers & Geology, vol. 7, Pergamon Press, Oxford (1990), p. 43-52
- (14) Method of interval kriging, Sciences de la Terre, Sér. Inf., no. 29 (1991), p. 17-33
- (15) Geostatistics in Bathymetry, Sciences de la Terre, Sér. Inf., no. 29 (1991), p. 35-46
- (16) HAMANN, I.M. and U.C. HERZFELD, On the effects of pre-analysis standardization, J. Geology, vol. 99, no. 4 (1991), p. 621-631
- (17) Quantitative spatial models of Atlantic primary productivity: An application of geomathematics, J. Geophys. Res., Oceans, vol. 97, no. C1 (1992), p. 717-732
- (18) Das Schwerefeld im Bereich des Explora Escarpments (Weddellmeer, Antarktis) - ein Vergleich zwischen Schiffsmessungen und Satellitendaten (GEOSAT), Zeitschrift f. Vermessungswesen, vol. 117, no. 8/9 (1992), p. 532-542

- (19) Least squares collocation, geophysical inverse theory, and geostatistics: A bird's eye view, *Geophys. J. Internat.*, vol. 111, no. 2 (1992), p. 237-249
- (20) MERRIAM, D.F., B.A. FUHR, and U.C. HERZFELD, An integrated approach to basin analysis and mineral exploration, in: Harff, J. and D.F. Merriam (eds.), *Computerized basin analysis for the prognosis of energy and mineral resources*, *Computers & Geology*, vol. 8, Pergamon Press, Oxford (1993), p. 197-214
- (21) ERIKSSON, M.G., H. BJÖRNSSON, U.C. HERZFELD, and P. HOLMLUND, The bottom topography of Storglaciären, *Stockholms Universitet Naturgeografiska Institutionen, Forskningsrapport 95*, ISSN 0346-7406, STOU-NG 95, Stockholm (1993), 48 p.
- (22) HERZFELD, U.C., I.I. KIM, J.A. ORCUTT, and C.G. FOX, Fractal geometry and seafloor topography: Theoretical concepts versus data analysis for the Juan de Fuca Ridge and the East Pacific Rise, *Annales Geophysicae*, vol. 11, no. 6 (1993), p. 532-541
- (23) DAVIS, J.C. and U.C. HERZFELD (eds.), *Computers in geology: 25 years of progress*, *Internat. Assoc. Math. Geol. Studies in Mathematical Geology*, no. 5, Oxford University Press, New York (1993), 298 p.
- (24) Fractals in geosciences - challenges and concerns, in: Davis, J.C. and U.C. Herzfeld (eds.), *Computers in geology: 25 years of progress*, *Internat. Assoc. Math. Geol. Studies in Mathematical Geology*, no. 5, Oxford University Press, New York (1993), p. 217-230
- (25) HERZFELD, U.C., C.S. LINGLE, and L. LEE, Geostatistical evaluation of satellite radar altimetry for high resolution mapping of Antarctic ice streams, *Annals Glaciol.*, vol. 17 (1993), p. 77-85
- (26) A method for seafloor classification using directional variograms, demonstrated for data from the western flank of the Mid-Atlantic ridge, *Math. Geol.*, vol. 25, no. 7 (1993), p. 901-924
- (27) HERZFELD, U.C., M.G. ERIKSSON, and P. HOLMLUND, On the influence of kriging parameters on the cartographic output - a study in mapping subglacial topography, *Math. Geol.*, vol. 25, no. 7 (1993), p. 881-900
- (28) MERRIAM, D.F., U.C. HERZFELD, and A. FÖRSTER, Integration of geophysical data for basin analysis, *Proc. Third Internat. Congress, Brazilian Geophys. Soc., Rio de Janeiro, 7-11 Nov. 1993*, (1993), vol. 2, p. 1025-1030.
- (29) HERZFELD, U.C. and A.L. BRODSCHOLL, On the geologic structure of the Explora Escarpment (Weddell Sea, Antarctica) revealed by satellite and shipboard data evaluation, *Marine Geophys. Res.*, vol. 16 (1994), p. 325-345
- (30) CERVENKA, P., U.C. HERZFELD, and C. DE MOUSTIER, Accuracy of the spatial representation of the seafloor with bathymetric sidescan sonars, *Marine Geophys. Res.*, vol. 16 (1994), p. 407-425
- (31) HERZFELD, U.C., C.S. LINGLE, and L. LEE, Recent advance of the grounding line of Lambert Glacier, Antarctica, deduced from satellite radar altimetry, *Annals Glaciol.*, vol. 20 (1994), p. 43-47
- (32) HERZFELD, U.C., I.I. KIM, and J.A. ORCUTT, Is the ocean floor a fractal? *Math. Geol.*, vol. 27, no. 3 (1995), p. 421-462
- (33) HERZFELD, U.C. and D.F. MERRIAM, Optimization techniques for integrating spatial data, *Math. Geol.*, vol. 27, no. 5 (1995), p. 559-588
- (34) HERZFELD, U.C. and C.A. HIGGINSON, Automated geostatistical seafloor classification — principles, parameters, feature vectors, and discrimination criteria, *Computers & Geosciences* vol. 22, no. 1 (1996), p. 35-52
- (35) HERZFELD, U.C. (guest editor), *Inverse Theory in the Earth Sciences*, Special Issue, *Math. Geol.*, vol. 28, no. 2 (1996), p. 133-251
- (36) Inverse theory in the Earth Science - an introductory overview with emphasis on Gandin's method of optimum interpolation, *Math. Geol.*, vol. 28, no. 2 (1996), p. 137-160

- (37) HERZFELD, U.C., B. KAUSCH, M. STAUBER, and A. THOMAS, Analysis of subscale ice-surface roughness from ultrasound measurements and its relevance for monitoring environmental changes from satellites, *Trierer Geograph. Studien*, 16 (1997, in press)
- (38) HERZFELD, U.C., C.S. LINGLE, C. FREEMAN, C.A. HIGGINSON, M.P. LAMBERT, L. LEE, and V. VORONINA, Monitoring changes of ice streams using time series of satellite-altimetry-based digital terrain models, *Math. Geol.*, vol. 29, no. 7 (1997, in press)
- (39) MERRIAM, D.F., U.C. HERZFELD, and B.A. FUHR, An integrated approach to petroleum exploration on the Pratt anticline, Pratt County, Kansas, *Amer. Assoc. Petrol. Geol. Bull.* (submitted)
- (40) HERZFELD, U.C. and H. MAYER, Surge of Bering Glacier and Bagley Ice Field, Alaska - an update of August 1995 and interpretation of brittle deformation patterns, *J. Glaciol.* (in press)
- (41) HERZFELD, U.C. and M.S. MATASSA, TRANSVIEW: A program for matching Universal Transverse Mercator (UTM) and geographic coordinates, *Computers & Geosciences* (submitted)
- (42) HERZFELD, U.C. and M.S. MATASSA, An atlas of Antarctica north of 72.1 °S from GEOSAT radar altimeter data, *Int. J. Remote Sensing* (in press)
- (43) Geostatistical interpolation and classification of remote-sensing data from ice surfaces, *Int. J. Remote Sensing* (submitted)